

# FLASHBlot™-SD Transfer Buffer


Rapid high efficiency semi-dry transfer buffer

**FLASHBlot™-SD Transfer Buffer** is designed for rapid semi-dry transfer of proteins from polyacrylamide gels (SDS-PAGE) to nitrocellulose or PVDF membranes using rapid semi-dry transfer systems. Transfer is compatible with commonly used detection methods such as membrane staining, chemiluminescent and fluorescent Western blotting.




## Advantages

- **FAST** – high ionic strength formulation allows for protein transfer in 3 to 10-minutes when used with a compatible high current semi-dry blotting system
- **COMPATIBLE** – use your existing high current semi-dry transfer apparatus
- **REPRODUCIBLE** – consistent transfer across entire blot
- **VERSATILE** – use nitrocellulose or PVDF membranes to achieve transfers with low background and high sensitivity with both chemiluminescent and fluorescent Western blots




**Figure 1. FLASHBlot™-SD is a versatile buffer, compatible with PVDF and Nitrocellulose membranes.** Chemiluminescent Western blot analysis of beta actin was performed on blots containing serially diluted HeLa lysate that was electrophoresed by SDS-PAGE then transferred to PVDF or Nitrocellulose membranes. Proteins were transferred from gel to membrane for 7 minutes at a constant current of 1.3 Amps.




**Figure 2. FLASHBlot™-SD produces fluorescent Western blots with low background and high sensitivity.** IR fluorescent Western blot analysis of phosphorylated STAT3 and GAPDH was performed on blots containing serially diluted HeLa lysate ( $\pm$ IFN $\alpha$  treatment) that was electrophoresed by SDS-PAGE then transferred to PVDF membranes. Proteins were transferred from gel to membrane for 7 minutes at a constant current of 1.3 Amps.

# FLASHBlot™-SD Transfer Buffer


Rapid high efficiency semi-dry transfer buffer



**Figure 3. FLASHBlot™-SD produces chemiluminescent Western blots with highest sensitivity.** Chemiluminescent Western blot analysis of hnRNP K and BAX was performed on blots containing serially diluted HeLa lysate that was electrophoresed by SDS-PAGE then transferred to nitrocellulose membranes. Proteins were transferred from gel to membrane for 7 minutes at a constant current of 1.3 Amps.



**Figure 4. FLASHBlot™-SD is compatible with various semi-dry blotters.** AdvanStain™ Iris membrane staining was performed on blots containing serially diluted *E.coli* lysate that was electrophoresed by SDS-PAGE then transferred to nitrocellulose membranes. Proteins were transferred from gel to membrane for 7 minutes at a constant current of 1.3 Amps with the Power Blotter and the Trans-Blot® Turbo™. Proteins were transferred from gel to membrane for 30 minutes at a constant current of 1.3 Amps with the Novex® semi-dry blotter.



**Figure 5. FLASHBlot™-SD outperforms competitive semi-dry transfer buffers.** (a) AdvanStain™ Total-PVDF fluorescent protein membrane staining was performed on blots containing serially diluted HeLa lysate that was electrophoresed by SDS-PAGE then transferred to PVDF membranes. Proteins were transferred from gel to membrane for 7 minutes at a constant current of 1.3 Amps. (b) After membrane staining was complete, an IR800 fluorescent Western blot analysis of GAPDH was performed. FLASHBlot-SD produced 4–8 fold higher Western blot sensitivity in comparison to other commercially available transfer buffers.

## Ordering Information

Catalog Number	Product	Size
R-03737-D50	FLASHBlot™-SD Transfer Buffer	500 ml
L-07125-120	Semi-Dry Blotting Paper (7 X 9cm), 10 Stacks	120 sheets
L-07126-120	Semi-Dry Blotting Paper (10 X 15cm) 10 Stacks	120 sheets



**Advansta Corporation**  
2140 Bering Drive | San Jose, CA 95131  
Tel: 650.325.1980 | Fax: 650.325.1904 | Email: [sales@advansta.com](mailto:sales@advansta.com)  
Product information: [www.advansta.com/products/FLASHBlot-SD-semi-dry-transfer-buffer](http://www.advansta.com/products/FLASHBlot-SD-semi-dry-transfer-buffer)

Copyright © 2025 Advansta. All rights reserved. The Advansta logo is a registered trademarks of the Company. AdvanStain™ and FLASHBlot™ are trademarks of the Company. TransBlot® Turbo™ are trademarks of Bio-Rad Laboratories. Novex® is a trademark of Thermo Fisher Scientific. All other trademarks, service marks and tradenames appearing in this brochure are the property of their respective owners.

[www.advansta.com](http://www.advansta.com)

